

Seeking Gauge Bileptons in Linear Colliders

Paul H. Frampton and Andrija Rašin

Department of Physics and Astronomy

University of North Carolina, Chapel Hill, NC 27599-3255

February 1, 2008

A promising direction to find physics beyond the standard model is to look for violation of $L_{e,\mu,\tau}$ conservation. In particular the process $e^-e^- \rightarrow \mu^-\mu^-$ with the exchange of a gauge bilepton has a striking signal without background and is predicted in the most economical model to have a cross-section an order of magnitude higher than previous estimates.

February 1, 2008

I. INTRODUCTION

Conservation laws, and their relation to symmetries, are fundamental to the theoretical physics of nuclei, condensed matter as well as high energy physics. This has already led to great advances and further progress will certainly come when the correct symmetries are better probed and identified.

It is expected that clear deviations from the Standard Model (SM) of particle physics should soon show up in experiment. Already the non-zero neutrino mass is one example of this - in the minimal SM neutrinos have no mass. Beyond that the best hope is to observe rare processes, especially those absolutely forbidden in the minimal SM. The best suited are processes which violate global conservation laws, since there is no reason to believe in their exactness. For example, baryon number (B) might be violated as in the proton decay predicted by *e.g.* grand unification. Even more amenable to test may be the processes which violate the separate lepton numbers $L_{e,\mu,\tau}$ since, even if $L = L_e + L_\mu + L_\tau$ is good, there is little reason to believe in $L_{e,\mu,\tau}$. Neutrino oscillations exemplify this.

One outstanding question of the SM is the occurrence of replicas of the quark-lepton families and one approach to that issue is the cancellation of chiral anomalies between asymmetric non-sequential families of the 331 model [1,2] (based on the gauge group $SU(3)_c \times SU(3)_L \times U(1)$). Recall that anomaly cancellation is crucial in many situations of model building beyond standard model, *e.g.* chiral color [3] and in string theory [4]. The 331 model simultaneously predicts that while L is a good quantum number perturbatively, processes do occur with $|\Delta L_i| = 2$ ($i = e, \mu, \tau$). There is no SM background, and so a process like $e^-e^- \rightarrow \mu^-\mu^-$ shown in Figure 1 provides a way to test [5] whether bileptonic Y^{--} gauge bosons are present, as they are if $SU(2)_L \subset SU(3)_L$ for the gauge group. Another model studied in this context is the SU(15) theory, where there are double charged gauge bileptons too, however with a smaller coupling (see




FIG. 1. Contribution of the gauge bilepton to the $e^-e^- \rightarrow \mu^-\mu^-$ scattering.

next Section). Among other theories which have doubly charged particles with $L = 2$ are the left-right models based on the group $SU(2)_L \times SU(2)_R \times U(1)_{B-L}$ [6]. However, the couplings of these states to the SM fermions are of Yukawa type and are not fixed. On the other hand, even though M_R , the scale of $SU(2)_R$ symmetry breaking, may be high, there are light double charged scalars and fermions [7,8], which may lie close to the weak scale. The situation also holds in the supersymmetric grand unified versions of left-right models, where complete double charged Higgs multiplets may be much lighter than the scales of left-right and grand unified symmetries [9]. The same experiments that will search for gauge bileptons will set important limits for these states as well [10].

In old work on the subject of gauge bileptons in the early 90's, the $L_{e,\mu}$ violating processes were calculated in the context of an SU(15) theory [11,12]. Here we point out that in the more economic and attractive 331 model, the cross-section $\sigma(e^-e^- \rightarrow \mu^-\mu^-)$ is larger by an order of magnitude than in SU(15). This result holds true in any model that embeds the gauge group $SU(2)_L$ in a single $SU(3)_L$.

The lower limit on $M(Y^{--})$ is currently claimed to be at 850 GeV [13] (however see discussion below, also see [14,15]), and with that value of $M(Y^{--})$ the cross-section for $\sigma(e^-e^- \rightarrow \mu^-\mu^-)$ is already ~ 5 fb at $\sqrt{s} = 100$ GeV and climbs very fast with s to several hundred fb at $\sqrt{s} =$

500 GeV. This implies a readily detectable event rate for an $L_{e,\mu}$ violating process in a collider with integrated luminosity of $10(\text{fb})^{-1}$ or higher, *e.g.* $10^{33}/\text{cm}^2\text{s}$ for a few months or $10^{34}/\text{cm}^2\text{s}$ for a week.

Some of the leading candidates presently discussed for the next generation of particle colliders (beyond LEP, Tevatron and LHC) are the Next Linear Collider (NLC), the (circular) Muon Collider (MC) and the Very Large Hadron Collider (VLHC). Of these the most advanced in design and preparation is the NLC. This is most often assumed to be an e^+e^- positron-electron collider with center of mass energy in the range above LEP: $200\text{GeV} < \sqrt{s} < 2000\text{GeV}$, although also being discussed is the desirability to operate e^+e^- at $\sqrt{s} = M_Z$ with high luminosity.

The motivation for such an NLC is to investigate physics in a presently-unexplored energy domain, although by the time of its start-up initial discoveries can be expected to have been made by LHC whereupon the NLC provides an excellent tool for detailed studies.

It is these machines that are ideally suited for finding the bilepton even for $\sqrt{s} < 200$ GeV (see Figure 2), but with luminosity higher than LEP. These machines could therefore initially run in both charge modes at such an energy.

II. CROSS SECTION

The interaction Lagrangian between the bilepton Y and leptons is given by

$$L_Y = \lambda Y_\mu^{++} e^T C \gamma^\mu P_- e + \lambda^\dagger Y_\mu^{--} \bar{e} \gamma^\mu P_- C \bar{e}^T \quad (1)$$

where λ is the 3×3 coupling matrix. For simplicity, in the following we will assume $\lambda^{ij} \sim \lambda \mathbf{1}$.

In the 331 model the group $SU(2)_L$ is completely embedded in the $SU(3)_L$ so that the gauge bilepton coupling is equal to the coupling of $SU(2)_L$

$$\lambda = \frac{g}{\sqrt{2}} \approx \frac{2.08e}{\sqrt{2}} \quad (2)$$

This coupling is larger than the coupling arising from the $SU(15)$ models where $\lambda \approx 1.19e/\sqrt{2}$ simply because the $SU(2)_L$ in that case is not residing entirely in a single $SU(3)_L$ but also partially in the $SU(6)_L$ subgroup of $SU(15)$ [16].

The differential cross section (see also [11,17]) for the gauge bilepton exchange in the process $e^-e^- \rightarrow \mu^-\mu^-$ is given by

$$\frac{d\sigma}{d\cos\theta} = \frac{\lambda^4}{32\pi} (1 + \cos^2\theta) \frac{s}{(s - m_Y^2)^2 + m_Y^2 \Gamma_Y^2} \quad (3)$$

cross sec in fb




FIG. 2. Total cross section in the 331 model for the $e^-e^- \rightarrow \mu^-\mu^-$ process as a function of bilepton mass at $\sqrt{s} = 100, 250, 500$ GeV.

where θ goes from 0 to $\pi/2$ reflecting the identical particles in the final states. Here m_Y and Γ_Y are the mass and decay width of the bilepton and s is the square of the center of mass energy. This leads to the total cross section

$$\sigma = \frac{\lambda^4}{24\pi} \frac{s}{(s - m_Y^2)^2 + m_Y^2 \Gamma_Y^2} \quad (4)$$

When the flavor and mass eigenbasis of leptons do not match, the results above must be corrected by $\lambda^2 \rightarrow \lambda^2 V_{ej} V_{je}^* V_{\mu j} V_{j\mu}^*$.

The total cross section as a function of the bilepton masses is plotted in Figure 2 for various values of $\sqrt{s}$ (We estimated the decay width of the bilepton as [17] $\Gamma_Y = \lambda^2 m_Y / 8\pi$.)

We see that already at $\sqrt{s} = 100$ GeV we get a cross section of about $\sigma \sim 5$ fb for a bilepton mass $m_Y = 850$ GeV, and for the same m_Y becomes 250 fb at $\sqrt{s} = 500$ GeV. This is about an *order of magnitude* higher than in the case of $SU(15)$ model.

Such a large cross section makes such a process highly visible. The situation is especially good for NLC-turned-into- e^-e^- -machine, even if it would run only at $\sqrt{s} = 100$ GeV, but with the desired luminosity of $10^{33}/\text{cm}^2\text{s}$.

This motivates a first run of future linear colliders (electron or muon) machines as *same charge lepton collider*. There are another two reasons for considering doing this: a) the additional cost to switch a machine from e^+e^- to e^-e^- is relatively negligible ; b) the highest energy that previously probed e^-e^- collisions was in 1971 at $\sqrt{s} = 1.12$ GeV [18], so this would present an opportunity to improve bounds on the bilepton mass considerably.

The spin 1 of the state Y can be checked by the angular distribution of Eq. (3). For spin 0 the distribution is isotropic.

The inverse process $\mu^+\mu^+ \rightarrow e^+e^+$ (μ^+ is easier to “cool” than μ^-) has the same cross section and could be probed at muon colliders, which are recently discussed as the next colliders in addition to e^+e^- machines.

III. BACKGROUNDS

Since the scattering violates L_e and L_μ it does not have any background in the standard model. However, a non-zero neutrino mass violates lepton number and one can easily find processes which involve neutrino mass that produce $e^-e^- \rightarrow \mu^-\mu^-$, as for example shown in Figure 3. Recently, first indications of neutrino mass have been found, and one can ask the question how serious a background the neutrino exchange can be to the bilepton exchange, given the mass ranges quoted in the atmospheric and solar neutrino data. In fact such calculations have been done recently [19] which find that the cross section is very small. The strongest bound for Figure 3a) comes from the neutrinoless double beta decay where it was found that the cross section for the process $e^-e^- \rightarrow WW$ is smaller than 2.5×10^{-7} fb for $\sqrt{s} = 100$ GeV and is thus quite negligible compared to the vector bilepton exchange. Bounds on contributions in Figure 3b) are found in [19] to be of a similar order, either because of a small light neutrino mass or a small mixing with the heavy Majorana neutrino masses. On the other hand, using fine tuning, very light right-handed neutrinos, or contributions from extra doubly charged scalar states, it is possible [19] to make the cross sections somewhat larger (typically $10^{-6} - 10^{-5}$ fb), but these are still much smaller than the bilepton signal. Also, we note that a cut would have to be applied for back-to-back muons in the final state, and therefore the signal to background ratio would be even higher.

In any case, non-back-to-back $\mu^-\mu^-$ in the final state are more likely to be from $e^-e^- \rightarrow Y^{--} \rightarrow \tau^-\tau^- \rightarrow \mu^-\mu^- + \text{neutrinos}$, which has the same cross-section as in Eq.(4), giving another positive signal for the Y^{--} .

A diagram not discussed in previous papers that contributes to the e^-e^- scattering and involves a pair of back-to-back scattered muons in the final states is shown in Figure 4. This process is not related directly to the neutrinoless double beta decay data. However, it is bound to be still smaller than the processes in Figure 3, because of the small neutrino masses. We estimate

$$T_{\text{Fig.4}} \sim \frac{1}{16\pi^2} |V_{ei}|^2 |V_{\mu j}|^2 \frac{m_{\nu_i} m_{\nu_j}}{M_W^4} \quad (5)$$

where V_{li} are the Kobayashi-Maskawa like mixing angles in the lepton sector. Even if we take typical neutrino




FIG. 3. Contributions from non-zero neutrino mass to background of $e^-e^- \rightarrow \mu^-\mu^-$.




FIG. 4. Another contribution to background of $e^-e^- \rightarrow \mu^-\mu^-$.

masses to be $\sim 1\text{eV}$ this process is *completely* negligible (down by over ten orders of magnitude) compared to the bilepton diagram (with m_Y in the TeV region) which has $T \sim V_{ei}V_{\mu j}^*/m_Y^2$.

IV. DISCUSSION

In this letter we have shown that the next generation of linear colliders, when run in the same charge mode, are especially suitable for search of doubly charged gauge bosons, that naturally appear in theories where $SU(2)_L$ is completely embedded in a new $SU(3)_L$ gauge group. Of special interest is the process $e^-e^- \rightarrow \mu^-\mu^-$ which has no background in the standard model. Even if neutrino masses are taken to be non-zero, with the current limits, the background to the bilepton exchange is completely negligible for $\sqrt{s} \leq 1$ TeV. This, in our opinion, strongly

motivates renewed interest in running the future colliders first as e^-e^- or $\mu^-\mu^-$ machines.

One aspect of the NLC, not usually mentioned in a physics paper but relevant to the real world and to our discussion, is its cost and the concomitant likelihood of being funded and constructed. For simplicity take the cost of an NLC to be linear in energy and assume it as one penny per eV. This estimate is very approximate, but, in any case, the minimization of E will clearly minimize the start-up cost of a machine which could then be extended to achieve higher E . The cost of running e^-e^- is perhaps slightly less than for e^+e^- , but both are needed for their physics interests. The e^+e^- mode at $\sqrt{s} = 100$ GeV provides useful machine physics and exploration of the Z pole at a higher luminosity than that of LEP ($5 \times 10^{31}/\text{cm}^2\text{s}$).

Finally, we comment on the current experimental limits on the bilepton mass. One of the strongest limits comes from the fermion pair production and lepton-flavor violating charged lepton decays [20] and is about 750 GeV. We see from Figure 2, that even for $\sqrt{s} = 100$ GeV we get huge signals for masses above the current limit and up to 1 TeV and well beyond. On the other hand an even stronger limit is claimed to come from the muonium-antimuonium conversion [13] which currently set the limit at about 850 GeV for the SU(15) like coupling. Since the diagram contributing to that process is exactly the same as the one contributing to the $e^-e^- \rightarrow \mu^-\mu^-$ scattering, one would at first sight think that the same limit applies, and so for a 331 coupling (which is larger than the SU(15)) the above limit would translate to an even stronger bound. However, there are several reasons why the bound from the muonium-antimuonium conversion does not have to apply in the same way. First, it was noted by Pleitez [14] that the limit of 850 GeV may be much less severe if there are additional diagrams with, for example, scalars contributing to the same process, that may cancel each other. In fact such extra scalars do appear in the minimal 331 model [1]. Second, the muonium-antimuonium conversion happens at energies which are much smaller than m_Y , typically less than 1 GeV. Then, however, for the $e^-e^- \rightarrow \mu^-\mu^-$ process which is happening at much higher energies (and with a nonnegligible center of energy s) the cancellation of diagrams may disappear and the process will be enhanced relatively to the muonium- antimuonium conversion.

Nevertheless, as we have shown, even for gauge bilepton masses in the TeV region, the signatures in $e^-e^- \rightarrow \mu^-\mu^-$ scattering would be significant at attainable design luminosities. In our opinion this warrants a serious examination of the possibility of including this mode in the designs of the future generation of linear colliders. It would provide a potentially very exciting experiment in the early stages of operation of such machines.

Acknowledgments

We thank David Burke for answering experimental queries and Ryan Rohm for a very useful discussion. This work was supported in part by the US Department of Energy under the Grant No. DE-FG02-97ER-41036.

-
- [1] P.H. Frampton, Phys. Rev. Lett. **69**, 2889 (1992).
 - [2] F. Pisano and V. Pleitez, Phys. Rev. **D46**, 410 (1992).
 - [3] P.H. Frampton and S.L. Glashow, Phys. Lett. **B190**, 157 (1987); Phys. Rev. Lett. **58**, 2168 (1987).
 - [4] P.H. Frampton and T.W. Kephart, Phys. Rev. Lett. **50**, 1343 (1983); *ibid*, **50**, 1347 (1983).
 - [5] See for example Proceedings to the 1st and 2nd Workshops (3rd to be published) on Electron-Electron Interactions on TeV Energies, ed. C.A. Heusch, Int. J. Mod. Phys. **A11**, 1523 (1996); *ibid* **A13**, 2217 (1998); See also <http://www.hep.psi.ch/e-e-.html>
 - [6] J. Pati and A. Salam, Phys. Rev. **D10**, 275 (1974); R. Mohapatra and J. Pati, Phys. Rev. **D11**, 2558 (1975); G. Senjanović and R. Mohapatra, Phys. Rev. **D12**, 1502 (1975); G. Senjanović, Nucl. Phys. **B153**, 334 (1979).
 - [7] C.S. Aulakh, A. Melfo and G. Senjanović, Phys. Rev. **D57**, 4174 (1998), hep-ph/9707256.
 - [8] See also Z. Chacko and R.N. Mohapatra, Phys. Rev. **D58**, 015001 (1998), hep-ph/9712359; C.S. Aulakh, A. Melfo, A. Rašin and G. Senjanović, Phys. Rev. **D58**, 115007 (1998), hep-ph/9712551; For a model with low M_R scale and broken R-parity see K. Huitu and J. Maalampi, Phys. Lett. **B344**, 217 (1995), hep-ph/9410342;
 - [9] C.S. Aulakh, B. Bajc, A. Melfo, A. Rašin and G. Senjanović, Phys. Lett. **B460**, 325 (1999), hep-ph/9904352.
 - [10] See, for example, J.F. Gunion, Int. J. Mod. Phys. **A13**, 2277 (1998), hep-ph/9803222; K. Huitu et al, hep-ph/9912405.
 - [11] P.H. Frampton and D. Ng, Phys. Rev. **D45**, 4240 (1992).
 - [12] P.H. Frampton, Mod. Phys. Lett. **A7**, 2017 (1992).
 - [13] L. Willmann *et al.*, Phys. Rev. Lett. **82**, 49 (1999); B. Ni *et al.*, Phys. Rev. **D48**, 1976 (1993).
 - [14] V. Pleitez, hep-ph/9905406.
 - [15] P. Das and P. Jain, hep-ph/9903432.
 - [16] P.H. Frampton and B.-H. Lee, Phys. Rev. Lett. **64**, 619 (1990).
 - [17] See also F. Cuyper and M. Raidal, Nucl. Phys. **B501**, 3 (1997), hep-ph/9704224. F. Cuyper and S. Davidson, Eur. Phys. J. **C2**, 503 (1998), hep-ph/9609487.
 - [18] W.C. Barber *et al*, Phys. Rev. **D3**, 2796 (1971).
 - [19] G. Belanger *et al*, Phys. Rev. **D53**, 6292 (1996), hep-ph/9508317. For various ways of somewhat relaxing these limits see P. Duka, J. Gluza and M. Zralek, Phys. Rev. **D58**, 053009 (1998), hep-ph/9804372. Also for an earlier attempt see C.A. Heusch and P. Minkowski, Nucl. Phys. **B416**, 3 (1994).
 - [20] M.B. Tully and G.C. Joshi, Phys. Lett. **B466**, 333 (1999), hep-ph/9905552.